



US005510245A

**United States Patent** [19]**Magers**[11] **Patent Number:** **5,510,245**[45] **Date of Patent:** **Apr. 23, 1996**[54] **COMPOSITION AND METHOD OF ASSAYING FOR KETONE BODIES**[75] Inventor: **Thomas A. Magers**, South Bend, Ind.[73] Assignee: **Bayer Corporation**, Elkhart, Ind.[21] Appl. No.: **182,405**[22] Filed: **Jan. 18, 1994****Related U.S. Application Data**

[62] Division of Ser. No. 941,707, Sep. 8, 1992, Pat. No. 5,326,697.

[51] Int. Cl.<sup>6</sup> ..... **C12Q 1/32; C12Q 1/26; C12Q 1/54; G01N 33/48**[52] U.S. Cl. .... **435/26; 435/25; 435/14; 435/4; 436/63; 436/74; 436/904**[58] Field of Search ..... **435/26, 25, 14, 435/4; 436/128, 904, 119, 63, 74**[56] **References Cited****U.S. PATENT DOCUMENTS**

3,528,888	9/1970	Deutsch	435/26
3,539,453	11/1970	Deutsch	435/26
3,867,258	2/1975	Forgione	435/26
4,758,323	7/1988	Davis et al.	435/26
4,873,260	10/1989	Alberts et al.	514/449
5,032,506	7/1991	Palmer et al.	435/26
5,126,247	6/1992	Palmer et al.	435/25

5,128,457	7/1992	Albarella et al.	436/119
5,187,105	2/1993	Albarella et al.	436/119
5,190,863	3/1993	Magers	435/26
5,326,697	7/1994	Magers	435/26

*Primary Examiner*—Douglas W. Robinson*Assistant Examiner*—Louise N. Leary*Attorney, Agent, or Firm*—Roger N. Coe[57] **ABSTRACT**

A composition, test device and method of determining the presence or concentration of ketone bodies, and specifically D- $\beta$ -hydroxybutyrate, in a test sample are disclosed. The test device includes a test pad comprising a suitable carrier matrix incorporating an indicator reagent composition capable of interacting with D- $\beta$ -hydroxybutyrate to produce a detectable or measurable response. In addition, a new and improved indicator reagent composition, comprising a) an indicator dye that is responsive to thiols, such as a substituted isobenzothiazolone, Ellman's reagent or a derivative of Ellman's reagent; b) D- $\beta$ -hydroxybutyrate dehydrogenase; c) lipoamide dehydrogenase; d) D,L-lipoamide; and e) nicotinamide adenine dinucleotide, is incorporated into the carrier matrix to provide an accurate and sensitive assay of a test sample for D- $\beta$ -hydroxybutyrate (DHBA) in particular, and for ketone bodies in general. The improved method and composition are especially useful in the assay of whole blood, blood serum, blood plasma and urine for ketone bodies.

**29 Claims, 14 Drawing Sheets****INT ADN IBTZ-I: LIGHT STRESS STUDY**  
**DAYLIGHT & U.V. COMBINED EXPOSURE**